

# The Geology And Vertebrate Paleontology Of Upper Eocene Strata In The Northeastern Part Of The Wind River Basin, Wyoming Part 2. The Mammalian Fauna Of The Badwater Area ( With 3 Plates)



[\[PDF\] Spacecraft \(Pointers\)](#)

[\[PDF\] Gluten Free Slow Cooker Recipes: Learn How to Prepare Delicious Gluten-Free Recipes with Your Slow Cooker](#)

[\[PDF\] Trauma Medico-Legal Medicine Anatomy and Surgery Volume 29 No. 6](#)

[\[PDF\] Revision Notes for the MCEM Part A \(Oxford Specialty Training: Revision Texts\)](#)

[\[PDF\] Complementary Health for Women: A Comprehensive Treatment Guide for Major Diseases and Common Conditions](#)

[\[PDF\] Tender Buttons](#)

[\[PDF\] \\*OP Sea of Shadows \(Wraith - the Oblivion\)](#)

**5 - PaleoPublications Home** 2. Perissodactyls from the Dry Creek and Badwater upper Eocene. 3. LEONTOLOGY OF UPPER EOCENE STRATA IN THE NORTHEASTERN PART OF THE WIND RIVER BASIN, WYOMING PART 2. THE MAMMALIAN FAUNA OF THE BADWATER AREA By C lewis GAZIN Curator, Division of Vertebrate Paleontology **View/Open - Smithsonian** (Palynology, Volume 2, 1978, pp. Geologic Map of the Southeastern Part of the Wind River Basin and Adjacent Areas in . The Geology And Vertebrate Paleontology Of Upper Eocene Strata In The Northeastern Part Of The Wind River Basin, Wyoming Part 2. The Mammalian Fauna Of The Badwater Area ( With 3 Plates). **The Geology And Vertebrate Paleontology Of Upper Eocene Strata** Halfway/Lapoint Fauna and regarded as the type fauna of the Duchesnean to about the upper three-quarters of the Halfway Duchesne River Formation faunas, wherein he the lower part of the Lapoint Member from road cuts . Institute of Technology vertebrate paleontological partial M1 or 2, SDSNH 130400 ?. **A REVISION OF TAPOCYON (CARNIVORAMORPHA), INCLUDING** the White River Formation and/or part of the Wind River Formation is shown 2. Vertebrates, Fossil. 3. Geology Wyoming Natrona. Co. I. Title. II. Series: about 750 feet of strata of the early Oligocene . the Flagstaff Rim area of this report (see geologic map, Figure 19) . So far as I can determine, no fossils of Eocene. Results 201 - 250 of 261 Your one-stop virtual bookshop dedicated to paleontology and related 2 in the Pacific War area. Institution Press, 1948, Miscellaneous Collections, 110 (2) : 1-3. . Geology and Vertebrate Paleontology of Upper Eocene Strata in the Wind River Basin, Wyoming. Part II. The Mammalian Fauna **The Geology And Vertebrate Paleontology Of Upper Eocene Strata** Berggren, W. A., 1986, Geochronology of the Eocene/Oligocene boundary, . Green River Basin, Wyoming: Journal of Vertebrate Paleontology, v. 15(3), p. of upper Eocene strata in the northeastern part of the Wind River Basin, Wyoming, Part 2. The mammalian fauna of the Badwater area: Smithsonian **MXieiiianJluseum - AMNH**

**Library Digital Repository - American** Fossiliferous strata of Hendry Ranch member of Tepee Trail formation. Ih. Figures. Page. 1. 23. Table. 1. Upper Eocene fossils from northeastern Wind River Basin, Wyoming. iii. 17

**GEOLOGY AND VERTEBRATE PALEONTOLOGY** Wind River. 1 Part 2 of this paper, The Mammalian Fauna of the Badwater Area, by. **Occurrence Detail 1039408589 - GBIF** Three new specimens of Tapocyon are described from Middle Eocene (Uintan NALMA) strata of San Diego County, California. The Geology and Vertebrate Paleontology of Upper Eocene Strata in the Northeastern Part of the Wind River Basin, Wyoming, Pt. 2, The Mammalian Fauna of the Badwater Area **Another Cretaceous Shark Tooth Id - Uk - Leaderboard - The Fossil** Results 451 - 498 of 498 Part III, - Geology and Vertebrate Paleontology of the Late Quaternary near Clovis New Mexico. Softbound, minor cover wear, text and plates in very good condition. Eocene Strata in the Northeastern Part of the Wind River Basin, Wyoming. Part II. The Mammalian Fauna of the Badwater Area. **GEOCHRONOLOGY AND MAGNETOSTRATIGRAPHY OF** Statistical data for upper and lower teeth of Amynodon advenus, Whistler Squat l.f. and Measurements and statistical data for M M 3 2, of Amynodon advenus and . Detailed descriptions of localities are on file at the Vertebrate Paleontology . Eocene beds on the north side of the Wind River Basin in the Badwater area. **Full Text (PDF) - SEPM Online Publications** Eocene Strata In. The Northeastern Part Of The Wind River Basin, Wyoming Part 2. The Mammalian Fauna Of The Badwater Area ( With 3 Plates) By C. Lewis. <http://findingaids/FARU7314.xml> **Charles Lewis** Prt. 2: The mammalian fauna of the badwater area. . vertebrate paleontology of upper Eocene strata in the Northeastern part of the Wind River Basin, Wyoming. **Full text of Eocene bad water - Internet Archive** The Mammalian Fauna Of The Badwater Area ( With 3 Plates) on Of Upper Eocene Strata In The Northeastern Part Of The Wind River Basin, Wyoming Part 2. **new records of small mammals from the middle eocene duchesne** Fossil vertebrates from the upper part of the Bridger Formation (Middle Eocene) at Tabernacle Butte in the northeastern part of the Green . and Dawson, 1966), near the northern edge of the Wind River Basin. strata are difficult to trace. 2 Although the late Wasatchian fauna from the Lost Cabin area has been used as **Journal of Paleontology - BioOne** The geology and vertebrate paleontology of upper Eocene strata in the northeastern part of the Wind River Basin, Wyoming. Part 2. The mammalian fauna of **Full text of The Geology and Vertebrate Paleontology of Upper** Eocene Strata In. The Northeastern Part Of The Wind River Basin, Wyoming Part 2. The Mammalian Fauna Of The Badwater Area ( With 3 Plates) By C. Lewis. **Report of the Workshop on Geologic Applications of Remote** Todd, J. Folios of the Geologic Atlas of the United States : Huron, South Dakota. . and Vertebrate Paleontology of Upper Eocene Strata in the Northeastern Part of the Wind River Basin, Wyoming. Part II. The Mammalian Fauna of the Badwater Area. Softbound, minor signs of use, text and plates in very good condition. **View/Open - Smithsonian Institution** ranged from vertebrate paleontology to geophysical modeling of continents. the Wind River/Bighorn Basin area of Wyoming in particular. 5. 2. Geologic Setting of Wind River/Bighorn Basin Area . . . . 7 e. 3. Remote Sensing in Sedimentary Basin Study . paleontology of upper Eocene strata in the northeastern part of. **The geology and vertebrate paleontology of upper Eocene strata in** Gazins research interests concern the vertebrate paleontology of the Tertiary and . Box 2 Folder 1 Chaffee, Robert Gibson, 1946, 1951, 1961 Folder 2 Cl-Co, Includes notes concerning fauna of the Badwater area, and correspondence with . Eocene Strata in the Northeastern part of the Wind River Basin, Wyoming. **View/Open - Smithsonian Institution** Eocene Strata In. The Northeastern Part Of The Wind River Basin, Wyoming Part 2. The Mammalian Fauna Of The Badwater Area ( With 3 Plates) By C. Lewis. - **The University of Texas at Austin** **The Geology And Vertebrate Paleontology Of Upper Eocene Strata** 1 Part I of this paper is a study of the geologic relations, in preparation by. Harry A. currence of upper Eocene on the north side of the Wind River Basin. Recognized by NO. 8. MAMMALIAN FAUNA, BADWATER AREAGAZIN. 3 helaletid (this is The discovery of Badwater vertebrate remains by Wood, Seton, and. **The Geology And Vertebrate Paleontology Of Upper Eocene Strata** Todd, J. Folios of the Geologic Atlas of the United States : Huron, South Dakota. . Printed wraps, xlib with all the usual signs, text and plates in good condition. . Vertebrate Paleontology of Upper Eocene Strata in the Northeastern Part of the Wind River Basin, Wyoming. Part II. The Mammalian Fauna of the Badwater Area. **The geology and vertebrate paleontology of upper Eocene strata in** Showing most informative content on 07/01/2014 in all areas . Contributions from the Museum of Paleontology, University of Michigan. Southeastern Geology, Vol.43, Number 3. . Journal of Mammalian Evolution, Vol.12, Numbers 1/2. Upper Eocene Strata in the Northeastern Part of the Wind River Basin, Wyoming. **275 - PaleoPublications** Department of Geology, Occidental College, Los Angeles, CA 90041 Dudnesnean land mammal ages are all middle Eocene age. vertebrate paleontologists (based on detailed stratigraphic . can 1, 2, 3, Torrejonian 1) and so on (Fig. 2). strata in the northeastern part of the Wind River Basin, Wyoming, Part 2.

**275 - PaleoPublications** The geology and vertebrate paleontology of Upper Eocene strata in the northeastern part of the Wind River Basin, Wyoming. Part 2. The mammalian fauna of the Badwater area. Smithsonian Miscellaneous and Archeology of Princeton College. bull. 3, art. iv: 46-53. Note: Wrong Plate number (Pl. V) listed in the text. **Structural Geology of the Wind River Basin, Wyoming - AbeBooks** Geology and vertebrate paleontology of upper Eocene strata in the northeastern part of the Wind River Basin, Wyoming. The geology and vertebrate **The Geology And Vertebrate Paleontology Of Upper Eocene Strata** Eocene Strata In. The Northeastern Part Of The Wind River Basin, Wyoming Part 2. The. Mammalian Fauna Of The Badwater Area ( With 3 Plates) By C. Lewis. **Next - PaleoPublications** 131, NO, 8 2 SMITHSONIAN MISCELLANEOUS COLLECTIONS VOL. The bone for the most part is a light buff color and the teeth amber to dark brown. . S MAMMALIAN FAUNA, BADWATER AREA GAZIN 7 Lack of uniformity of opinion . One of the more important discoveries in the upper Eocene of the Wind River **275 - PaleoPublications Home** Three new specimens of Tapocyon are described from Middle Eocene (Uintan NALMA) strata of San Diego County, California. The Geology and Vertebrate Paleontology of Upper Eocene Strata in the Northeastern Part of the Wind River Basin, Wyoming, Pt. 2, The Mammalian Fauna of the Badwater Area